

Metoidioplasty without UL **Postoperative Instructions**

1. No Strenuous physical activity of any type during the first 6 weeks (No bending, pushing, pulling, lifting (greater than 20lbs) straining or excessive walking)
2. Resume your regular diet as tolerated (see soft diet suggestions below)
3. Avoid excessive alcohol intake
4. No smoking for at least one month following surgery
5. Use pain medications as needed for pain/discomfort and please take antibiotics as directed
6. Swelling is completely normal, and the phallus and scrotum will often swell to 2-3 times in size after the compression dressing is removed at the first postoperative appointment. You can apply circumferential pressure in a downward motion to help with the swelling after the first week.

Suggested Supplies

- **Hand mirror**
 - Allows the patient to view surgical area while changing dressings
- **Aquaphor or Neosporin**
 - May provide relief and comfort where new skin is exposed
- **Thick pillow or Medical donut**
 - For comfort while sitting or travelling
- **Ice packs or Bags of frozen vegetable**
 - Moldable ice packs or bags of frozen vegetables provide temporary pain relief and decrease swelling
- **Extra pairs of underwear**
 - Most patients prefer underwear that is looser and seamless if possible. Tight underwear can be uncomfortable and rub against the newly exposed skin. Seams can also cause discomfort.
- **Sweat pants and loose jeans**
 - For comfort and to accommodate the catheter with drainage bag and drain that will still be present postoperatively.
- **Cushiony cup or pad for inside your underwear**
 - Past patients have purchased soft cups from Andrew Christian. A heavily padded cup from a bra will also suffice. As patients have transitioned to wearing tighter briefs postoperatively, this provided comfort.
- **Lap desk**
 - This may allow you to accomplish tasks and serve as protection against dogs/pets/children on your lap.
- **Extra Strength Tylenol (Acetaminophen)**
 - Transition to taking Tylenol after your pain medication for mild to moderate pain relief. Do not take Tylenol concurrently with your pain medication as the pain medication also contains acetaminophen and large amounts may cause liver damage.
- **Maxi Pads (super absorbent)**
 - Absorb any drainage after dressings are removed.

Diet

BROWNSTEIN & CRANE SURGICAL SERVICES

- Resume your normal diet.
- Lack of activity postoperatively and pain medications can contribute to constipation. Helpful preventative measures including taking:
 - Prune Juice
 - Miralax may be used in the event extra help is needed
 - Avoid Metamucil or other fiber drinks as they may contribute to more constipation during the period of reduced activity

Important Contact Information:

****For medical or logistical questions, please phone the office during normal business hours****

Office Phone: 415.625.3230

Office Hours: Monday - Friday, 9:00 am- 5:00 pm

For any of the below situations, please phone the office during normal business hours with any questions.

****Swelling** is a normal postoperative occurrence; you can use ice to decrease this (20 min on/20 off)

****Postoperative pain** is a normal postoperative occurrence; please take the medication as prescribed

******As of October 6, 2014, postoperative pain medications can no longer be called into pharmacies. Please contact our office to arrange pick up of a prescription if you anticipate needing additional medications. Medications will not be called in over the weekend.******

****Nausea** is a normal side effect of postoperative pain medication; if you experience nausea, please stop taking the prescription pain medication and switch to Extra Strength Tylenol or contact our office for an antiemetic prescription.

****Constipation** is a normal side effect of postoperative pain medication; you may take an over the counter laxative or stool softener.

****When the office is closed and there is a medical question that requires a same day response, please call Ursula at 415-378-9061**

For any emergencies please call 9-1-1.

If any of the situations listed below occur outside of business hours, please contact Dr. Crane or Chen on his cell phone ONLY in the following events:

****Fever over 101° F**

****Pus or drainage from the incision sites coupled with a foul odor**

Dr. Crane's Cell Phone: 603-667-5555

Dr. Chen's Cell Phone: 412-390-9513